



Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



MATHEMATICS

9709/03

Paper 3 Pure Mathematics 3

For examination from 2020

SPECIMEN PAPER

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **20** pages. Blank pages are indicated.

- 2 (a) Expand $(1 + 3x)^{-\frac{1}{3}}$ in ascending powers of x , up to and including the term in x^2 , simplifying the coefficients. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) State the set of values of x for which the expansion is valid. [1]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

3 (a) Sketch the graph of $y = |2x - 3|$.

[1]

(b) Solve the inequality $3x - 1 > |2x - 3|$.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Verify by calculation that this equation has a root between 1 and 2.

[2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Use the iterative formula $a_{n+1} = \frac{1}{2}(3 - \ln a_n)$ to calculate a correct to 2 decimal places, showing the result of each iteration to 4 decimal places.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 5 (a) Show that $\frac{d}{dx}(x - \tan^{-1}x) = \frac{x^2}{1+x^2}$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Show that $\int_0^{\sqrt{3}} x \tan^{-1}x \, dx = \frac{2}{3}\pi - \frac{1}{2}\sqrt{3}$. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

A series of 24 horizontal dotted lines for writing.

6 The complex numbers $1 + 3i$ and $4 + 2i$ are denoted by u and v respectively.

(a) Find $\frac{u}{v}$ in the form $x + iy$, where x and y are real. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) State the argument of $\frac{u}{v}$. [1]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

In an Argand diagram, with origin O , the points A , B and C represent the complex numbers u , v and $u - v$ respectively.

- (c) State fully the geometrical relationship between OC and BA . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (d) Show that angle $AOB = \frac{1}{4}\pi$ radians. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

.....



In the diagram, $OABC$ is a pyramid in which $OA = 2$ units, $OB = 4$ units and $OC = 2$ units. The edge OC is vertical, the base OAB is horizontal and angle $AOB = 90^\circ$. Unit vectors \mathbf{i} , \mathbf{j} and \mathbf{k} are parallel to OA , OB and OC respectively. The midpoints of AB and BC are M and N respectively.

- (a) Express the vectors \vec{ON} and \vec{CM} in terms of \mathbf{i} , \mathbf{j} and \mathbf{k} . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Calculate the angle between the directions of \vec{ON} and \vec{CM} . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) Show that the length of the perpendicular from M to ON is $\frac{3}{5}\sqrt{5}$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

9



The diagram shows the curve $y = \sin^2 2x \cos x$ for $0 \leq x \leq \frac{1}{2}\pi$, and its maximum point M .

(a) Find the x -coordinate of M .

[6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Using the substitution $u = \sin x$, find the area of the shaded region bounded by the curve and the x -axis. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 10 In a chemical reaction, a compound X is formed from two compounds Y and Z .

The masses in grams of X , Y and Z present at time t seconds after the start of the reaction are x , $10 - x$ and $20 - x$ respectively. At any time the rate of formation of X is proportional to the product of the masses of Y and Z present at the time. When $t = 0$, $x = 0$ and $\frac{dx}{dt} = 2$.

- (a) Show that x and t satisfy the differential equation

$$\frac{dx}{dt} = 0.01(10 - x)(20 - x). \quad [1]$$

.....

.....

.....

.....

.....

.....

- (b) Solve this differential equation and obtain an expression for x in terms of t . [9]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) State what happens to the value of x when t becomes large. [1]

.....

.....

.....

.....

.....

