When is your child too sick for school?


School attendance is a powerful predictor of school success and on-time graduation. Research shows that when students miss fewer than 10% of their school days, they have much stronger success in school. Use these tips to tell when your child is too sick to attend school.

Send your child to school if they...



Don't feel great but are able to participate in normal activities.


Keep your child home if they...



Feel sick and have a fever of 100.4°F or higher, test positive for flu or COVID-19, or can't participate in normal activities.

Have thrown up in the past 24 hours.





Have diarrhea and increased stool frequency.

Have a rash of unknown origin. They can return to school when a healthcare provider says it's okay.






Have an oozing sore or wound you can't cover or contain.


Call a healthcare provider if they...



Have a fever of 100.4°F or higher for more than 2 days.


Throw up or have diarrhea for more than 2 days.





Have the sniffles for more than a week and aren't getting better.

Have asthma symptoms after using asthma medicine. If they have trouble breathing after using an inhaler, call 911.





This is general guidance. Schools, healthcare providers, or the Health Department may advise differently based on individual circumstances.

Washington Office of Superintendent of
PUBLIC INSTRUCTION

Adapted from Tacoma-Pierce County Health Department in partnership with the WA State Department of Health

¿Cuándo está su hijo demasiado enfermo para ir a la escuela?

La asistencia a la escuela es un gran factor predictivo del éxito escolar y de la graduación a tiempo. Las investigaciones muestran que los estudiantes, cuando faltan menos del 10 % de sus días escolares, tienen un éxito mucho mayor en la escuela. Aproveche estos consejos para saber cuándo su hijo está demasiado enfermo para asistir a la escuela.

Envíe a su hijo a la escuela si...



No se siente muy bien, pero puede participar en actividades normales.

Mantenga a su hijo en casa si...



Se siente enfermo y tiene fiebre de 100.4 °F o más, da positivo en la prueba de gripe o COVID-19, o no puede participar en actividades normales.


Ha vomitado en las últimas 24 horas.





Tiene diarrea y una mayor frecuencia de las deposiciones.

Tiene una erupción de origen desconocido. Puede regresar a la escuela cuando un proveedor de atención médica lo autorice.





Tiene una llaga o herida supurante que no puede cubrir ni contener.


Llame a un proveedor de atención médica si...



Tiene fiebre de 100.4 °F o más durante más de 2 días.


Vomita o tiene diarrea durante más de 2 días.





Lleva más de una semana con congestión nasal y no mejora.

Tiene síntomas de asma después de usar medicamentos para el asma. Si tiene problemas para respirar después de usar un inhalador, llame al 911.





Esta es una guía general. Las escuelas, los proveedores de atención médica o el Health Department pueden ofrecer consejos diferentes según las circunstancias individuales.

Washington Office of Superintendent of
PUBLIC INSTRUCTION

Adaptado del Tacoma-Pierce County Health Department en asociación con el Washington (WA) State Department of Health.